## Chronic Inflammatory Demyelinating Polyneuropathy

Symmetric proximal and distal weakness, length dependent loss of large fiber sensation (Typical)

Sensory Predominant

**Motor Predominant** 

Chronic Immune Sensory Polyneuropathy

Acute onset CIDP

Focal CIDP (Restricted to one limb)

Distal Acquired Demyelinating Symmetric Neuropathy (DADS)\* Lewis Sumner
Syndrome/
Multifocal
Acquired
Demyelinating
Sensory and Motor
neuropathy
(MADSAM)

AĬĎP

- Nadir within 4 weeks since onset
- Fluctuating GBS has <2 relapses and does not have any relapse after 8 weeks

**SIDP** 

• Nadir in 4-8 weeks since onset

CIDP

- Nadir after 8 weeks since onset
- Acute onset CIDP has  $\geq 3$  relapses (some after 8 weeks of symptom onset) or can be chronically progressive

AIDP – Acute Inflammatory Demyelinating Polyneuropathy

SIDP – Subacute Inflammatory Demyelinating Polyneuropathy

CIDP – Chronic Inflammatory Demyelinating Polyneuropathy

GBS – Guillain-Barre Syndrome

\*If diagnosed as DADS, check IgM antibodies. If positive, it should be diagnosed as a paraproteinemic neuropathy and not CIDP. 50-70% of DADS are paraproteinemic neuropathies

Hariharan Venkataraman, Final year MBBS, JSSMC, India