TARDIVE DISORDERS

Classic tardive dyskinesia	Oral-buccal lingual stereotypy +/- trunk and limb involvement (usually distal)
Tardive dystonia	Focal, segmental or generalized dystonia; more common in young men; blepharospasm, retrocollis, opisthotonus, arm extension and pronation w/ wrist flexion.
Tardive akathisia	An internal sensation of restlessness with the inability to remain still
Tardive myoclonus	Spontaneous or stimulus-sensitive jerk-like movements; UE>LE, ↑ w/ sustained posture and/or voluntary movement
Tardive tics	Adult-onset motor/phonic tics
Tardive tremor	Postural and/or resting tremor, 3–5Hz
Tardive chorea	Random, jerk-like movements that migrate from one muscle group to the next
Tardive gait	"Dance-like": short steps on toes followed by longer strides "Duck-like": wide-based, unsteady gait with short stride length and steppage pattern
Tardive Parkinsonism	Rest tremor, bradykinesia, and rigidity, which appear after DA receptor blocker discontinuation and last for months-years
Withdrawal emergent syndrome	Generalized choreiform movements mainly of the neck, trunk, and limbs in children after DA receptor blocker discontinuation; self-limited
Tardive ocular deviation	Painless conjugate eye deviations; Fixed stare → conjugate spasmodic movements of the eyes, generally upward
Tardive pain	Neuropathic-type pain in the oral and/or genital regions

THOMAS PIEKUT

Based on: Savitt and Jankovic, 2018 (doi: 10.1016/j.jns.2018.02.005)